Evolutionary Ecology, 1995, **9**, 678–684

Index to Volume 9 of Evolutionary Ecology

SUBJECT INDEX

3–4 year vole; parental effort; reversed-size dimorphism; owl; 30
abundance; geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; biogeography; fossils; 586
Adelges japonicus; intergall migration; evolution of dispersal; ESS model; gall-forming aphid; 542
adult lifespan; birds; mammals; invariants; 288
aggregation; polymorphism; *Drosophila*; patches; soft-selection; 508
agonistic encounters; nest failure; cost of fighting; burying beetles; communal breeding; co-operation; 125
alevins; functional response; Atlantic salmon; *Salmo salar*; timing of emergence; predation; fry; 411
allelic sensitivity; genetic correlation; phenotypic plasticity; gene regulation; 154
allometric scaling; convergent evolution; ecomorphology; bird migration; 520
alternative tactics; ESS model; limited breeding site; host density; anemonefish; sex change; 292
anemonefish; sex change; alternative tactics; ESS model; limited breeding site; host density; 292
angiosperm diversification; comparative biology; vertebrate dispersal; fruit dispersal; seed dispersal; 93
annual fluctuations; mictotine rodents; *Microtus agrestis*; population cycles; seasonality; temporary habitats; 64
antiherbivore defences; co-existence; compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; 605
Aphididae; intraclonal variability; predator escape behaviour; maternal effects; artificial selection; Homoptera; 397
Apis mellifera; *Brassica campestris*; isolation by distance; maximum likelihood; pollen deposition; 304
Aquarius remigis; sexual size dimorphism; body size; food limitation; foraging; mating; trade-off; Gerridae; 369
archipelagos; endemism; islands; skinks; speciation; species diversity; Pacific Ocean; 529
artificial selection; Homoptera; Aphididae; intraclonal variability; predator escape behaviour; maternal effects; 397
Asobara tabida; *Leptopilina boulardi*; parasitoid specificty; geographical variation; encapsulation ability; *Drosophila melanogaster*; parasitoid; 10
astigmatic mites; genotype-by-environment interaction; threshold concept; dispersal traits; unpredicatably varying habitats; 341
Atlantic salmon; *Salmo salar*; timing of emergence; predation; fry; alevins; functional response; 411
attractiveness; pollination; selfing rate; size-number trade-off; optimal allocation model; seed size; 495
avoidable and unavoidable mortality; density-dependent mortality; perturbation; life-history; evolution; 478

bees; compound eye; insect vision; flower form; 236
biogeography; fossils; abundance; geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; 586
biological control; population stability; co-evolution; parasitoid; population dynamics; refuge; 633
biomechanics; convergence; hydrophily; seagrass; pollination; submarine pollination; 139
bird migration; allometric scaling; convergent evolution; ecomorphology; 520
birds; mammals; invariants; adult lifespan; 288
body size; food limitation; foraging; mating; trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; 369
bottleneck; inbreeding depression; predator-prey models; extinction; immigration; 1
Brassica campestris; isolation by distance; maximum likelihood; pollen deposition; *Apis mellifera*; 304
burying beetles; communal breeding; cooperation; agonistic encounters; nest failure; cost of fighting; 125

circadian rhythm; emergence; insect parasitoid; locomotor activity; protandry; 467
clines; colonization; *D. subobscura*; *D. obscura*; quantitative traits; 453
clonal disintegration; evolutionary context; plant clonal integration; plant physiological integration; clonal evolution; 575
clonal evolution; clonal disintegration; evolutionary context; plant clonal integration; plant physiological integration; 575
co-evolution; parasitoid; population dynamics; refuge; biological control; population stability; 633
co-existence; compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; 605
colonization; *D. subobscura*; *D. obscura*; quantitative traits; clines; 453
communal breeding; co-operation; agonistic encounters; nest failure; cost of fighting; burying beetles; 125
comparative biology; vertebrate dispersal; fruit dispersal; seed dispersal; angiosperm diversification; 93
compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; co-existence; 605
competition; density dependence; fish communities; *Salmo salar*; *Salvelinus fontinalis*; habitat selection; 169
compound eye; insect vision; flower form; bees; 236
convergence; hydrophily; seagrass; pollination; submarine pollination; biomechanics; 139
convergent evolution; ecomorphology; bird migration; allometric scaling; 520
co-operation; agonistic encounters; nest failure; cost of fighting; burying beetles; communal breeding; 125

Evolutionary Ecology, 1995, **9**, 678–684

Index to Volume 9 of Evolutionary Ecology

SUBJECT INDEX

3–4 year vole; parental effort; reversed-size dimorphism; owl; 30
abundance; geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; biogeography; fossils; 586
Adelges japonicus; intergall migration; evolution of dispersal; ESS model; gall-forming aphid; 542
adult lifespan; birds; mammals; invariants; 288
aggregation; polymorphism; *Drosophila*; patches; soft-selection; 508
agonistic encounters; nest failure; cost of fighting; burying beetles; communal breeding; co-operation; 125
alevins; functional response; Atlantic salmon; *Salmo salar*; timing of emergence; predation; fry; 411
allelic sensitivity; genetic correlation; phenotypic plasticity; gene regulation; 154
allometric scaling; convergent evolution; ecomorphology; bird migration; 520
alternative tactics; ESS model; limited breeding site; host density; anemonefish; sex change; 292
anemonefish; sex change; alternative tactics; ESS model; limited breeding site; host density; 292
angiosperm diversification; comparative biology; vertebrate dispersal; fruit dispersal; seed dispersal; 93
annual fluctuations; mictotine rodents; *Microtus agrestis*; population cycles; seasonality; temporary habitats; 64
antiherbivore defences; co-existence; compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; 605
Aphididae; intraclonal variability; predator escape behaviour; maternal effects; artificial selection; Homoptera; 397
Apis mellifera; *Brassica campestris*; isolation by distance; maximum likelihood; pollen deposition; 304
Aquarius remigis; sexual size dimorphism; body size; food limitation; foraging; mating; trade-off; Gerridae; 369
archipelagos; endemism; islands; skinks; speciation; species diversity; Pacific Ocean; 529
artificial selection; Homoptera; Aphididae; intraclonal variability; predator escape behaviour; maternal effects; 397
Asobara tabida; *Leptopilina boulardi*; parasitoid specificty; geographical variation; encapsulation ability; *Drosophila melanogaster*; parasitoid; 10
astigmatic mites; genotype-by-environment interaction; threshold concept; dispersal traits; unpredicatably varying habitats; 341
Atlantic salmon; *Salmo salar*; timing of emergence; predation; fry; alevins; functional response; 411
attractiveness; pollination; selfing rate; size-number trade-off; optimal allocation model; seed size; 495
avoidable and unavoidable mortality; density-dependent mortality; perturbation; life-history; evolution; 478

bees; compound eye; insect vision; flower form; 236
biogeography; fossils; abundance; geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; 586
biological control; population stability; co-evolution; parasitoid; population dynamics; refuge; 633
biomechanics; convergence; hydrophily; seagrass; pollination; submarine pollination; 139
bird migration; allometric scaling; convergent evolution; ecomorphology; 520
birds; mammals; invariants; adult lifespan; 288
body size; food limitation; foraging; mating; trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; 369
bottleneck; inbreeding depression; predator-prey models; extinction; immigration; 1
Brassica campestris; isolation by distance; maximum likelihood; pollen deposition; *Apis mellifera*; 304
burying beetles; communal breeding; cooperation; agonistic encounters; nest failure; cost of fighting; 125

circadian rhythm; emergence; insect parasitoid; locomotor activity; protandry; 467
clines; colonization; *D. subobscura*; *D. obscura*; quantitative traits; 453
clonal disintegration; evolutionary context; plant clonal integration; plant physiological integration; clonal evolution; 575
clonal evolution; clonal disintegration; evolutionary context; plant clonal integration; plant physiological integration; 575
co-evolution; parasitoid; population dynamics; refuge; biological control; population stability; 633
co-existence; compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; 605
colonization; *D. subobscura*; *D. obscura*; quantitative traits; clines; 453
communal breeding; co-operation; agonistic encounters; nest failure; cost of fighting; burying beetles; 125
comparative biology; vertebrate dispersal; fruit dispersal; seed dispersal; angiosperm diversification; 93
compensatory feeding; ESS; herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; co-existence; 605
competition; density dependence; fish communities; *Salmo salar*; *Salvelinus fontinalis*; habitat selection; 169
compound eye; insect vision; flower form; bees; 236
convergence; hydrophily; seagrass; pollination; submarine pollination; biomechanics; 139
convergent evolution; ecomorphology; bird migration; allometric scaling; 520
co-operation; agonistic encounters; nest failure; cost of fighting; burying beetles; communal breeding; 125

cost of fighting; burying beetles; communal breeding; co-operation; agonistic encounters; nest failure; 125
cost of reproduction; stochastic demography; fitness; life-history; red deer; selection pressures; 559

D. obscura; quantitative traits; clines; colonization; *D. subobscura*; 453
D. subobscura; *D. obscura*; quantitative traits; clines; colonization; 453
demography; dispersal; ecological niche; evolution; heterogenous environments; natural selection; source-sink populations; 38
density dependence; fish communities; *Salmo salar*; *Salvelinus fontinalis*; habitat selection; competition; 169
density-dependent mortality; perturbation; life-history; evolution; avoidable and unavoidable mortality; 478
diet choice; static information; obligate information; facultative information; 18
dispersal; ecological niche; evolution; heterogenous environments; natural selection; source-sink populations; demography; 38
dispersal traits; unpredicatably varying habitats; astigmatic mites; genotype-by-environment interaction; threshold concept; 341
Drosera tracyi; natural selection; pollination; resource levels; 382
Drosophila; patches; soft-selection; aggregation; polymorphism; 508
Drosophila melanogaster; parasitoid; *Asobara tabida*; *Leptopilina boulardi*; parasitoid specificty; geographical variation; encapsulation ability; 10
dynamic programming; parasitoid behaviour; host feeding; dynamic state variable models; 217
dynamic programming; feather quality; seasonality; survival rate; reproductive success; molt pattern; 354
dynamic state variable models; dynamic programming; parasitoid behaviour; host feeding; 217

ecological niche; evolution; heterogenous environments; natural selection; source-sink populations; demography; dispersal; 38
ecomorphology; bird migration; allometric scaling; convergent evolution; 520
emergence; insect parasitoid; locomotor activity; protandry; circadian rhythm; 467
encapsulation ability; *Drosophila melanogaster*; parasitoid; *Asobara tabida*; *Leptopilina boulardi*; parasitoid specificty; geographical variation; 10
endemism; islands; skinks; speciation; species diversity; Pacific Ocean; archipelagos; 529
Erythoxylum havanense; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; 204
ESS; herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; co-existence; compensatory feeding; 605
ESS model; gall-forming aphid; *Adelges japonicus*; intergall migration; evolution of dispersal; 542
ESS model; limited breeding site; host density; anemonefish; sex change; alternative tactics; 292
evolution; avoidable and unavoidable mortality; density-dependent mortality; perturbation; life-history; 478
evolution; heterogenous environments; natural selection; source-sink populations; demography; dispersal; ecological niche; 38
evolution of dispersal; ESS model; gall-forming aphid; *Adelges japonicus*; intergall migration; 542
evolutionarily stable strategy; exploitive competition; phenology; life cycle; renewable resource; 328
evolutionary context; plant clonal integration; plant physiological integration; clonal evolution; clonal disintegration; 575
evolutionary stable strategies; kin selection; fluctuating environments; geometric mean fitness; 185
exploitive competition; phenology; life cycle; renewable resource; evolutionarily stable strategy; 328
extinction; immigration; bottleneck; inbreeding depression; predator-prey models; 1
extinction; population extremes; Hurst exponents; year-to-year variability; 429

facultative information; diet choice; static information; obligate information; 18
feather quality; seasonality; survival rate; reproductive success; molt pattern; dynamic programming; 354
fig wasps; sex ratio; morph rate; sexual selection; 550
fish communities; *Salmo salar*; *Salvelinus fontinalis*; habitat selection; competition; density dependence; 169
fitness; life-history; red deer; selection pressures; cost of reproduction; stochastic demography; 559
fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; Erythoxylum havanense; 204
fitness set; life-history evolution; phenotypic plasticity; spatial heterogeneity; optimal reaction norm; 242
flower form; bees; compound eye; insect vision; 236
flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; 204
flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; 204
flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-

dependent selection; tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; 204
fluctuating environments; geometric mean fitness; evolutionary stable strategies; kin selection; 185
food; foraging time; lemming; mammal; microtine rodent; optimization; predation; seasonality; risk; reproduction; vole; 45
food limitation; foraging; mating; trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; body size; 369
foraging; mating; trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; body size; food limitation; 369
foraging time; lemming; mammal; microtine rodent; optimization; predation; seasonality; risk; reproduction; vole; food; 45
forest Macrolepidoptera; phylogenetic analysis; phenology; population dynamics; reduced wings; 275
fossils; abundance; geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; biogeography; 586
frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; 204
frozen niche-variation model; growth; survival and reproduction; maintenance of sex; unisexual reproduction; 258
fruit dispersal; seed dispersal; angiosperm diversification; comparative biology; vertebrate dispersal; 93
fry; alevins; functional response; Atlantic salmon; *Salmo salar*; timing of emergence; predation; 411
functional response; pollinator foraging behaviour; pollen limitation; plant regulation; 421
functional response; Atlantic salmon; *Salmo salar*; timing of emergence; predation; fry; alevins; 411

gall-forming aphid; *Adelges japonicus*; intergall migration; evolution of dispersal; ESS model; 542
gap formation; resource allocation model; sapling growth; 444
gene regulation; allelic sensitivity; genetic correlation; phenotypic plasticity; 154
genetic correlation; phenotypic plasticity; gene regulation; allelic sensitivity; 154
genotype-by-environment interaction; threshold concept; dispersal traits; unpredicatably varying habitats; astigmatic mites; 341
geographic range; turnover; mollusc assemblages; Pleistocene; macroecology; biogeography; fossils; abundance; 586
geographical variation; encapsulation ability; *Drosophila melanogaster*; parasitoid; *Asobara tabida*; Leptopilina boulardi; parasitoid specificty; 10
geometric mean fitness; evolutionary stable strategies; kin selection; fluctuating environments; 185
Gerridae; *Aquarius remigis*; sexual size dimorphism; body size; food limitation; foraging; mating; trade-off; 369
growth; survival and reproduction; maintenance of sex; unisexual reproduction; frozen niche-variation model; 258

habitat selection; competition; density dependence; fish communities; *Salmo salar*; *Salvelinus fontinalis*; 169
haplodiploid; inclusive fitness; one-locus genetic model; sex ratio; sibmating; inbreeding depression; 318
herbivore mobility; indirect interactions; nutritive quality; antiherbivore defences; co-existence; compensatory feeding; ESS; 605
heterogenous environments; natural selection; source-sink populations; demography; dispersal; ecological niche; evolution; 38
Homoptera; Aphididae; intraclonal variability; predator escape behaviour; maternal effects; artificial selection; 397
host density; anemonefish; sex change; alternative tactics; ESS model; limited breeding site; 292
host feeding; dynamic state variable models; dynamic programming; parasitoid behaviour; 217
host range; host specialization; trade-offs; performance; normal versus novel hosts; host specificity; 82
host specialization; trade-offs; performance; normal versus novel hosts; host specificity; host range; 82
host specificity; host range; host specialization; trade-offs; performance; normal versus novel hosts; 82
Hurst exponents; year-to-year variability; extinction; population extremes; 429
hydrophily; seagrass; pollination; submarine pollination; biomechanics; convergence; 139

immigration; bottleneck; inbreeding depression; predator-prey models; extinction; 1
inbreeding depression; haplodiploid; inclusive fitness; one-locus genetic model; sex ratio; sibmating; 318
inbreeding depression; predator-prey models; extinction; immigration; bottleneck; 1
inclusive fitness; one-locus genetic model; sex ratio; sibmating; inbreeding depression; haplodiploid; 318
indirect interactions; nutritive quality; antiherbivore defences; co-existence; compensatory feeding; ESS; herbivore mobility; 605
insect parasitoid; locomotor activity; protandry; circadian rhythm; emergence; 467
insect vision; flower form; bees; compound eye; 236
intergall migration; evolution of dispersal; ESS model; gall-forming aphid; *Adelges japonicus*; 542
intraclonal variability; predator escape behaviour; maternal effects; artificial selection; Homoptera; Aphididae; 397
invariants; adult lifespan; birds; mammals; 288
islands; skinks; speciation; species diversity; Pacific Ocean; archipelagos; endemism; 529
isolation by distance; maximum likelihood; pollen deposition; *Apis mellifera*; *Brassica campestris*; 304

kin selection; fluctuating environments; geometric mean fitness; evolutionary stable strategies; 185

lemming; mammal; microtine rodent; optimization; predation; seasonality; risk; reproduction; vole; food; foraging time; 45

Leptopilina boulardi; parasitoid specificty; geographical variation; encapsulation ability; *Drosophila melanogaster*; parasitoid; *Asobara tabida*; 10
life cycle; renewable resource; evolutionarily stable strategy; exploitive competition; phenology; 328
life-history; evolution; avoidable and unavoidable mortality; density-dependent mortality; perturbation; 478
life-history; red deer; selection pressures; cost of reproduction; stochastic demography; fitness; 559
life-history evolution; phenotypic plasticity; spatial heterogeneity; optimal reaction norm; fitness set; 242
limited breeding site; host density; anemonefish; sex change; alternative tactics; ESS model; 292
litter size; trade-off; offspring size; offspring number; 57
locomotor activity; protandry; circadian rhythm; emergence; insect parasitoid; 467

macroecology; biogeography; fossils; abundance; geographic range; turnover; mollusc assemblages; Pleistocene; 586
maintenance of sex; unisexual reproduction; frozen niche-variation model; growth; survival and reproduction; 258
male strategies; mating competition; mating systems; paternal care; sexual selection; 662
mammal; microtine rodent; optimization; predation; seasonality; risk; reproduction; vole; food; foraging time; lemming; 45
mammals; invariants; adult lifespan; birds; 288
maternal effects; artificial selection; Homoptera; Aphididae; intraclonal variability; predator escape behaviour; 397
mating; trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; body size; food limitation; foraging; 369
mating competition; mating systems; paternal care; sexual selection; male strategies; 662
mating systems; paternal care; sexual selection; male strategies; mating competition; 662
maximum likelihood; pollen deposition; *Apis mellifera*; *Brassica campestris*; isolation by distance; 304
mean crowding; resource-defence polygyny; run-away evolution; non-adaptive aggregation; sexual selection; 118
microtine rodent; optimization; predation; seasonality; risk; reproduction; vole; food; foraging time; lemming; mammal; 45
Microtus agrestis; population cycles; seasonality; temporary habitats; annual fluctuations; mictotine rodents; 64
mictotine rodents; *Microtus agrestis*; population cycles; seasonality; temporary habitats; annual fluctuations; 64
mollusc assemblages; Pleistocene; macroecology; biogeography; fossils; abundance; geographic range; turnover; 586
molt pattern; dynamic programming; feather quality; seasonality; survival rate; reproductive success; 354
morph rate; sexual selection; fig wasps; sex ratio; 550

natural selection; pollination; resource levels; *Drosera tracyi*; 382
natural selection; source-sink populations; demography; dispersal; ecological niche; evolution; heterogenous environments; 38
nest failure; cost of fighting; burying beetles; communal breeding; cooperation; agonistic encounters; 125
non-adaptive aggregation; sexual selection; mean crowding; resource-defence polygyny; run-away evolution; 118
normal versus novel hosts; host specificity; host range; host specialization; trade-offs; performance; 82
nutritive quality; antiherbivore defences; co-existence; compensatory feeding; ESS; herbivore mobility; indirect interactions; 605

obligate information; facultative information; diet choice; static information; 18
offspring number; litter size; trade-off; offspring size; 57
offspring size; offspring number; litter size; trade-off; 57
one-locus genetic model; sex ratio; sibmating; inbreeding depression; haplodiploid; inclusive fitness; 318
optimal allocation model; seed size; attractiveness; pollination; selfing rate; size-number trade-off; 495
optimal reaction norm; fitness set; life-history evolution; phenotypic plasticity; spatial heterogeneity; 242
optimization; predation; seasonality; risk; reproduction; vole; food; foraging time; lemming; mammal; microtine rodent; 45
owl; 3–4 year vole; parental effort; reversed-size dimorphism; 30

Pacific Ocean; archipelagos; endemism; islands; skinks; speciation; species diversity; 529
parasitoid; *Asobara tabida*; *Leptopilina boulardi*; parasitoid specificty; geographical variation; encapsulation ability; *Drosophila melanogaster*; 10
parasitoid; population dynamics; refuge; biological control; population stability; co-evolution; 633
parasitoid behaviour; host feeding; dynamic state variable models; dynamic programming; 217
parasitoid specificty; geographical variation; encapsulation ability; *Drosophila melanogaster*; parasitoid; *Asobara tabida*; *Leptopilina boulardi*; 10
parental effort; reversed-size dimorphism; owl; 3–4 year vole; 30
patches; soft-selection; aggregation; polymorphism; *Drosophila*; 508
paternal care; sexual selection; male strategies; mating competition; mating systems; 662
performance; normal versus novel hosts; host specificity; host range; host specialization; trade-offs; 82
perturbation; life-history; evolution; avoidable and unavoidable mortality; density-dependent mortality; 478
phenology; life cycle; renewable resource; evolutionarily stable strategy; exploitive competition; 328
phenology; population dynamics; reduced wings; forest Macrolepidoptera; phylogenetic analysis; 275
phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; 204

phenotypic plasticity; gene regulation; allelic sensitivity; genetic correlation; 154
phenotypic plasticity; spatial heterogeneity; optimal reaction norm; fitness set; life-history evolution; 242
phylogenetic analysis; phenology; population dynamics; reduced wings; forest Macrolepidoptera; 275
plant clonal integration; plant physiological integration; clonal evolution; clonal disintegration; evolutionary context; 575
plant physiological integration; clonal evolution; clonal disintegration; evolutionary context; plant clonal integration; 575
plant regulation; functional response; pollinator foraging behaviour; pollen limitation; 421
Pleistocene; macroecology; biogeography; fossils; abundance; geographic range; turnover; mollusc assemblages; 586
pollen deposition; *Apis mellifera*; *Brassica campestris*; isolation by distance; maximum likelihood; 304
pollen limitation; plant regulation; functional response; pollinator foraging behaviour; 421
pollination; resource levels; *Drosera tracyi*; natural selection; 382
pollination; submarine pollination; biomechanics; convergence; hydrophily; seagrass; 139
pollination; selfing rate; size-number trade-off; optimal allocation model; seed size; attractiveness; 495
pollinator foraging behaviour; pollen limitation; plant regulation; functional response; 421
polymorphism; *Drosophila*; patches; soft-selection; aggregation; 508
population cycles; seasonality; temporary habitats; annual fluctuations; mictotine rodents; *Microtus agrestis*; 64
population dynamics; refuge; biological control; population stability; co-evolution; parasitoid; 633
population dynamics; reduced wings; forest Macrolepidoptera; phylogenetic analysis; phenology; 275
population extremes; Hurst exponents; year-to-year variability; extinction; 429
population stability; co-evolution; parasitoid; population dynamics; refuge; biological control; 633
predation; fry; alevins; functional response; Atlantic salmon; *Salmo salar*; timing of emergence; 411
predation; seasonality; risk; reproduction; vole; food; foraging time; lemming; mammal; microtine rodent; optimization; 45
predator escape behaviour; maternal effects; artificial selection; Homoptera; Aphididae; intraclonal variability; 397
predator-prey models; extinction; immigration; bottleneck; inbreeding depression; 1
protandry; circadian rhythm; emergence; insect parasitoid; locomotor activity; 467

quantitative traits; clines; colonization; *D. subobscura*; *D. obscura*; 453

red deer; selection pressures; cost of reproduction; stochastic demography; fitness; life-history; 559
reduced wings; forest Macrolepidoptera; phylogenetic analysis; phenology; population dynamics; 275
refuge; biological control; population stability; co-evolution; parasitoid; population dynamics; 633
renewable resource; evolutionarily stable strategy; exploitive competition; phenology; life cycle; 328
reproduction; vole; food; foraging time; lemming; mammal; microtine rodent; optimization; predation; seasonality; risk; 45
reproductive success; molt pattern; dynamic programming; feather quality; seasonality; survival rate; 354
resource allocation model; sapling growth; gap formation; 444
resource levels; *Drosera tracyi*; natural selection; pollination; 382
resource-defence polygyny; run-away evolution; non-adaptive aggregation; sexual selection; mean crowding; 118
reversed-size dimorphism; owl; 3–4 year vole; parental effort; 30
risk; reproduction; vole; food; foraging time; lemming; mammal; microtine rodent; optimization; predation; seasonality; 45
run-away evolution; non-adaptive aggregation; sexual selection; mean crowding; resource-defence polygyny; 118

Salmo salar; *Salvelinus fontinalis*; habitat selection; competition; density dependence; fish communities; 169
Salmo salar; timing of emergence; predation; fry; alevins; functional response; Atlantic salmon; 411
Salvelinus fontinalis; habitat selection; competition; density dependence; fish communities; *Salmo salar*; 169
sapling growth; gap formation; resource allocation model; 444
seagrass; pollination; submarine pollination; biomechanics; convergence; hydrophily; 139
seasonality; risk; reproduction; vole; food; foraging time; lemming; mammal; microtine rodent; optimization; predation; 45
seasonality; survival rate; reproductive success; molt pattern; dynamic programming; feather quality; 354
seasonality; temporary habitats; annual fluctuations; mictotine rodents; *Microtus agrestis*; population cycles; 64
seed dispersal; angiosperm diversification; comparative biology; vertebrate dispersal; fruit dispersal; 93
seed size; attractiveness; pollination; selfing rate; size-number trade-off; optimal allocation model; 495
selection pressures; cost of reproduction; stochastic demography; fitness; life-history; red deer; 559
selfing rate; size-number trade-off; optimal allocation model; seed size; attractiveness; pollination; 495
sex change; alternative tactics; ESS model; limited breeding site; host density; anemonefish; 292
sex ratio; sibmating; inbreeding depression; haplodiploid; inclusive fitness; one-locus genetic model; 318
sex ratio; morph rate; sexual selection; fig wasps; 550
sexual selection; male strategies; mating competition; mating systems; paternal care; 662
sexual selection; fig wasps; sex ratio; morph rate; 550
sexual selection; mean crowding; resource-defence polygyny; run-away evolution; non-adaptive aggregation; 118
sexual size dimorphism; body size; food limitation; foraging; mating; trade-off; Gerridae; *Aquarius remigis*; 369
sibmating; inbreeding depression; haplodiploid; inclusive fitness; one-locus genetic model; sex ratio; 318
size-number trade-off; optimal allocation model; seed size; attractiveness; pollination; selfing rate; 495

skinks; speciation; species diversity; Pacific Ocean; archipelagos; endemism; islands; 529
soft-selection; aggregation; polymorphism; *Drosophila*; patches; 508
source-sink populations; demography; dispersal; ecological niche; evolution; heterogenous environments; natural selection; 38
spatial heterogeneity; optimal reaction norm; fitness set; life-history evolution; phenotypic plasticity; 242
speciation; species diversity; Pacific Ocean; archipelagos; endemism; islands; skinks; 529
species diversity; Pacific Ocean; archipelagos; endemism; islands; skinks; speciation; 529
static information; obligate information; facultative information; diet choice; 18
stochastic demography; fitness; life-history; red deer; selection pressures; cost of reproduction; 559
submarine pollination; biomechanics; convergence; hydrophily; seagrass; pollination; 139
survival and reproduction; maintenance of sex; unisexual reproduction; frozen niche-variation model; growth; 258
survival rate; reproductive success; molt pattern; dynamic programming; feather quality; seasonality; 354

temporal variation in selection; frequency-dependent selection; tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; 204
temporary habitats; annual fluctuations; mictotine rodents; *Microtus agrestis*; population cycles; seasonality; 64
threshold concept; dispersal traits; unpredicatably varying habitats; astigmatic mites; genotype-by-environment interaction; 341
timing of emergence; predation; fry; alevins; functional response; Atlantic salmon; *Salmo salar*; 411
trade-off; Gerridae; *Aquarius remigis*; sexual size dimorphism; body size; food limitation; foraging; mating; 369
trade-off; offspring size; offspring number; litter size; 57
trade-offs; performance; normal versus novel hosts; host specificity; host range; host specialization; 82
tropical dry forest; watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; 204
turnover; mollusc assemblages; Pleistocene; macroecology; biogeography; fossils; abundance; geographic range; 586

unisexual reproduction; frozen niche-variation model; growth; survival and reproduction; maintenance of sex; 258
unpredicatably varying habitats; astigmatic mites; genotype-by-environment interaction; threshold concept; dispersal traits; 341

vertebrate dispersal; fruit dispersal; seed dispersal; angiosperm diversification; comparative biology; 93
vole; food; foraging time; lemming; mammal; microtine rodent; optimization; predation; seasonality; risk; reproduction; 45

watering experiments; flowering induction; *Erythoxylum havanense*; fitness; flowering synchrony; flowering phenology; phenotypic natural selection; temporal variation in selection; frequency-dependent selection; tropical dry forest; 204

year-to-year variability; extinction; population extremes; Hurst exponents; 429

AUTHOR INDEX

Ackerman, J. D. , 139
Adler, G. H. , 529
Adler, F. R. , 304
Andrade, M. C. B. , 397
Ariño, A. , 429
Augner, M. , 605
Austin, C. A. , 529

Benton, T. G. , 559
Blanckenhorn, W. U. , 369
Boulétreau, F. , 467
Brännäs, E. , 411
Brown, L. P. , 57
Brown, J. H. , 586

Charnov, E. L. , 57, 288, 662
CluttonBrock, T. H. , 559
Collier, T. R. , 217

Dafni, A. , 236
De Meeûs, T. , 131
Denver, K. , 318
Dirzo, R. , 204
Dominguez, C. A. , 204
Downhower, J. F. , 57
Dudley, R. , 529
Dytham, C. , 508

Enquist, B. J. , 586
Ezoe, H. , 328

Fairbairn, D. J. , 369
Fouillet, F. , 467

Grant, A. , 559
Greeff, J. M. , 550
Guillén, A. , 453

Haefner, J. W. , 1
Hakkarainen, H. , 30
Hartt, L. , 1
Hattori, A. , 292
Hawkes, K. , 662
Hedenström, A. , 354
Hernandez, M-J. , 478
Hochberg, M. E. , 131, 633
Holmgren, N. , 354
Holt, R. D. , 633
Hunter, A. F. , 275

Ingvarsson, P. K. , 421

Jordan, M. A. , 586
Joshi, A. , 82

Kawecki, T. J. , 38
Kelly, C. K. , 575
Kevan, P. G. , 236
Knülle, W. , 341
Korpimäki, E. , 30
Kotler, B. P. , 18
Kraaijeveld, A. R. , 10

León, M-J. , 478
Lomnicki, A. , 64
Lundberg, S. , 421
Lundberg, P. , 45

Mangel, M. , 304
Mazer, S. J. , 93
McNamara, J. M. , 185
Mehrhoff, L. A. , 617
Mitchell, W. A. , 18
Mönkkönen, M. , 520
Morris, W. F. , 304

Oksanen, L. , 45
Ozaki, K. , 542
Papaceit, M. , 453
Pegueroles, G. , 453
Pigliucci, M. , 154
Pimm, S. , 429
Pompanon, F. , 467
Prevosti, A. , 453
Preziosi, R. F. , 369

Quintana, A. , 453

Renaud, F. , 131
Rodriguez, M. A. , 169
Rogers, A. R. , 662
Roitberg, B. D. , 397

Sakai, S. , 444, 495
Schlichting, C. D. , 154
Serra, L. , 453
Shorrocks, B. , 508
Sibly, R. M. , 242

Taylor, P. D. , 318
Thompson, J. N. , 82
Tiffney, B. H. , 93
Trumbo, S. T. , 125
Turkington, R. , 617

van Alphen, J. J. M. , 10

Wade, M. J. , 118
Weeks, S. C. , 258
Wilson, P. , 382

Yamamura, N. , 292